

Immunological Non Responders to HAART in Botswana: Characteristics & Risk factors

Gaone Retshabile

Background

High global burden of HIV/AIDS

- 34 million people living with HIV/AIDS worldwide
- 23 million in Sub-Saharan Africa
- ↑ prevalence Southern Africa (UNAIDS,2012)

Botswana prevalence rate 25% adults

- 316 000 adults living with HIV
- 15 000 children under 14 years (National AIDS Council, 2010)

Background

Usually untreated infection → immunological deterioration

- CD4 T-helper cell decay, ↑ viral load (McMichael *et al.*, 2010; Moir, Tae-Wook & Fauci, 2011)

Immune activation

- Cytokine activation (inflammatory)
 - GALT damage → microbial translocation
 - AICD → immune exhaustion
- (Douek *et al.*, 2009; McMichael *et al.*, 2010)

Background


HAART

- ↓ morbidity & mortality, ↑ CD4 count, ↓ viral load (Bofill *et al.*, 2006, Bussmann *et al.*, 2008)
- >500 cells/ μ l = non HIV positive morbidity & mortality (Gaardbo *et al.*, 2012; Nixon & Landay, 2010).

Response heterogeneity

- Poor immunologic response despite virologic control
- 15-30% <200 cell/ μ l by 12 months treatment
- clinical risk (Casotti *et al.*, 2011; Gaardbo *et al.*, 2012; Nakanjako *et al.*, 2011)

Background



Gaardbo *et al.*, 2012

Justification

Conflicting data over INR & OI/death risk association

(Nakanjako et al., 2008; Gazolla *et al.*, 2009; Hermans *et al.*, 2010)

- HR AIDS/death (<200 vs. >350) 10 and CD4 <200 vs. 200-350 at 8.5 (Casotti et al., 2011)

Late treatment initiation (Bussmann *et al.*, 2008; Wester *et al.*, 2009; Esposito *et al.*, 2011)

- 48% VCT diagnosis <200 cells/ μ l (UNAIDS, 2012)
- \downarrow CD4 in HIV –ve in Botswana (Bussmann *et al.*, 2004)

Treatment strategies

- IL-2, IL-7 (Gaardbo *et al.*, 2012; Gazolla *et al.*, 2009)

Hypothesis

Immunological non response to HAART is associated with increased risk for the development of opportunistic infections and or death

Main Objective

Main objective

- To characterise patient factors associated with immunological non response to HAART in Botswana and its association with the risk of developing opportunistic infection and or death

Specific Objectives

Specific objectives

- Determine frequency of INR at 12 months ART
- Characterise polyfunctionality of activated PBMC
- Determine plasma levels of LPS, sCD14
- Determine systemic cytokine profiles of participants
- Associate baseline characteristics, immunological response, PBMC responses, microbial translocation and cytokine profiles with frequency of OI and or death

Methods

Population

- >18 yrs, consented, ART outcome studies
- Botswana MOH clearance, HSPH

Inclusion Criteria

- <200 cells/ μ l CD4 initiation, VL <400 copies/ml by 6 months, clinical data

Exclusion criteria

- 2 consecutive VL >400 copies/ml post 6 months

Methods

Definition: <200 cells/ μ l by 12 months ART

Data abstraction

- CD4 counts, Viral load, OI frequencies

Determine frequency INR

Laboratory analysis

- ELISPOT for PBMC responses
- ELISA (LPS, sCD14, cytokine profiles)

References

- Bussmann, H, Wester, CW, Masupu, KV, Peter, T, Gaolekwe, SM, Kim, S, Reich, AM, Ahn, S, Wu, Y, Thior, I, Essex, M and Marlink, R 2004, 'Low CD4⁺ T-Lymphocyte values in Human Immunodeficiency Virus-Negative Adults in Botswana', *Clinical and Diagnostic Laboratory Immunology*, vol. 11, no. 5, pp. 930-935.
- Bussmann, H, Wester, CW, Ndwapu, N, Grundmann, N, Gaolathe, T, Puvimanasinghe, J, Avalos, A, Mine, M, Seipone, K, Essex, M, DeGruttola, V and Marlink, RG 2008, 'Five Year Outcomes of Initial Patients Treated in Botswana's National Antiretroviral Treatment Programme', *AIDS*, vol. 22, no. 17, pp. 2303-2311.
- Casotti, JA, Passos, LN, Oliveira, FJP and Cerrutti jr., C 2011, 'Factors associated with paradoxical immune response to antiretroviral therapy in HIV infected patients: a case control study', *BMC Infectious Diseases*, vo. 11, pp. 306-316.
- Douek, DC, Roederer, M and Koup, RA 2009, Emerging 'Concepts in the Immunopathogenesis of AIDS', *Annual Review of Medicine*, vol. 60, pp. 471-484.
- Gaardbo, JC, Hartling, HJ, Gerstoft, J and Nielsen, SD 2012, 'Incomplete Immune Recovery in HIV: Mechanisms, Relevance for Clinical Care, and Possible Solutions', *Clinical and Developmental Immunology*, vol. 2012, no. 670957, pp. 1-17.
- Gazolla, L, Tincati, C, Bellistri, GM, d'Arminio Monforte, A and Marchetti, G 2009, 'The Absence of CD4⁺ T-cell Recovery Despite Receipt of Virologically Suppressive Highly Active Antiretroviral Therapy: Clinical Risk, Immunological Gaps and Therapeutic Options', *Clinical Infectious Diseases*, vol. 48, no. 3, pp. 328-337.

References

- Hermans, SB, Kiragga, AN, Schaefer, P, Kambugu, A, Hoepelman, AIM and Manabe YC 2010, 'Incident Tuberculosis during Antiretroviral Therapy Contributes to Suboptimal Immune Reconstitution in a Large Urban Clinic in Sub-Saharan Africa', *PloS One*, vol. 5, no 5, pp. 1-8.
- McMichael, AJ, Borrow, P, Tomaras, GD, Goonetilleke, N and Haynes, BF 2010, 'The immune response during acute HIV-1 infection: clues for vaccine development', *Nature Reviews Immunology*, vol. 10, no 1, pp. 11-23.
- Moir, S, Tae-Wook, C & Fauci, AS 2011, 'Pathogenic Mechanisms of HIV Disease', *Annual Review of Pathology: Mechanism of Disease*, vol. 11, no 6, pp. 223-248.
- Nakanjako, D, Ssewanyana, I, Mayanja-Kizza, H, Kiragga, A, Colebunders, R, Manabe, YC, Nabatanzi, R, Kanya, MR and Cao, H 2011, 'High T-cell immune activation and immune activation among individuals with suboptimal CD4 recovery after 4 years of antiretroviral therapy in an African cohort', *BMC Infectious Diseases*, vol. 11, pp. 43-51.
- National AIDS Coordinating Agency, 2010, *Botswana Country Report: 2008-2009*, Gaborone: National AIDS Coordinating Agency.
- UNAIDS, 2012, *Epidemic update and health sector progress towards Universal Access: 2011*, Geneva: United Nations Joint Programme on HIV/AIDS.